

COVID-19

EHC TEMPORARY PROTOCOL - LEVONORGESTREL 1500MG SUPPLY

This protocol is being introduced for the duration of the CV19 pandemic. It is recognised that face to face consultations should be suspended until further notice. This simplified process is not clinical best practice, but will allow supply of EHC after a brief conversation at a safe social distance OR after a telephone consultation. Ulipristal should not be supplied.

CLIENT IDENTIFIER: DATE:

CLIENT D.O.B. FACE TO FACE OR PHONE?

HAVE YOU HAD UNPROTECTED SEXUAL INTERCOURSE IN THE LAST 3 DAYS (72 HOURS)
YES NO (CLIENT NOT SUITABLE)

WHAT MEDICATION DO YOU NORMALLY TAKE?

ANY OF THESE ENZYME INDUCERS? IF SO – DOUBLE DOSE!

WHAT IS YOUR WEIGHT? WEIGHT OVER 70kg? GIVE DOUBLE DOSE!

HOW MANY LEVONORGESTREL SUPPLIED?
ONE TWO

PHARMACIST NAME:

ENZYME INDUCING DRUGS:

- Antifungals: griseofulvin.
- Antibiotics: rifampicin and rifabutin.
- Anti-epileptics: carbamazepine, eslicarbazepine, phenytoin, phenobarbital, primidone, oxcarbazepine, topiramate.
- Central nervous system stimulant: modafinil.
- Antiretroviral drugs: nelfinavir, nevirapine, ritonavir.
- St John's wort.